

Providence St. Joseph Health

## Providence St. Joseph Health Digital Commons

---

Providence Pharmacy PGY1 Program at Providence Portland and Providence St. Vincent Medical Centers 2021

Providence Pharmacy PGY1 Program at Providence Portland and Providence St. Vincent Medical Centers

---

5-2021

### Evaluation and Implementation of a New Phenobarbital Protocol for Alcohol Withdrawal Management in the Emergency Department

Sarah J. Kim

Sara Clark

Joshua L. Floyd

Follow this and additional works at: [https://digitalcommons.psjhealth.org/oaa\\_ppmcstvin\\_21](https://digitalcommons.psjhealth.org/oaa_ppmcstvin_21)



Part of the [Emergency Medicine Commons](#), and the [Pharmacy and Pharmaceutical Sciences Commons](#)

---



# Evaluation and Implementation of a New Phenobarbital Protocol for Alcohol Withdrawal Management in the Emergency Department

Sarah J. Kim, PharmD, Sara M. Clark, PharmD, Joshua L. Floyd, PharmD

## Background

- Alcohol withdrawal syndrome can lead to substantial health care costs and mortality.
- Although historically benzodiazepines (BZDs) are considered the mainstay treatment for alcohol withdrawal, some studies have shown that phenobarbital may be the safer and preferred alternative option.
- Advantages of phenobarbital are its mechanism of action, more predictable pharmacokinetics, and greater therapeutic index.
- A prospective study by Rosenson et al. 2012 found that a single 10 mg/kg IV dose of phenobarbital versus standardized lorazepam protocol resulted in decreased ICU admissions (8% vs. 25%).
- A retrospective study by Tidewell et al. 2018 demonstrated shorter hospital stays, less incidence of mechanical ventilation and less use of adjunctive agents with phenobarbital compared to that of BZDs.
- The phenobarbital protocol was approved by the Oregon Region P&T in October 2020.

## Purpose

- The purpose is to evaluate the safety and efficacy of phenobarbital use compared to BZDs use for the management of alcohol withdrawal in the emergency department (ED) at two large tertiary medical centers.

## Objectives

- Evaluate the safety and efficacy of phenobarbital in alcohol withdrawal compared to BZDs.
- Discuss the new phenobarbital protocol that was implemented in this study.
- Assess protocol compliance and areas of improvement.

## Methodology

- Institutional Review Board (IRB)-approved
- Electronic health record (EHR)-based retrospective chart review of patients in the ED at two large tertiary medical centers
- Study population:
  - Patients  $\geq 18$  years admitted to the ED with the primary diagnosis of 'alcohol withdrawal' with a presenting CIWA/PAWSS scores and received phenobarbital or BZD therapy (based on CIWA scores ranging from 9 to  $\geq 25$ )
  - Patients were stratified according to presenting CIWA/PAWSS scores (see protocol for complete scale)
- Study period:
  - Jan 1, 2020– Mar 31, 2021
- Included first 60 patients that met inclusion criteria between determined study period
- Exclusion criteria:
  - Allergy or hypersensitivity to phenobarbital or BZDs
  - Pregnancy
  - Documented history of acute intermittent porphyria, hepatic encephalopathy
  - Use of phenobarbital prior to admission as an anti-epileptic
  - Concomitant HIV medication use contraindicated with phenobarbital (Evotaz, Prezcoib, Tivicay, Doravirine, Genvoya, Stribild)
- Primary Outcome: ICU admission for alcohol withdrawal from the ED
- Secondary Outcomes: Incidence of mechanical ventilation, use of restraints, use of adjunctive agents, and protocol compliance
- Evaluation of phenobarbital protocol compliance based on three factors: appropriate CIWA/PAWSS risk stratification, correct weight for loading dose and met safety parameters

## Preliminary Results

Table 1. Patient Baseline Characteristics

Characteristic	Phenobarbital Group (n=30)	BZD Group (n=30)
Age, Years (Mean)	47 (range 25-75)	45 (range 29-70)
Sex, Male, No. (%)	20 (66.7)	19 (63)
Weight, Kg (Mean)	79.2 (range 45.4-117.9)	82.4 (range 48.5-113.4)
<b>Race, No. (%)</b>		
White or Caucasian	26 (86.7)	27 (90)
Black or African American	2 (6.7)	0
Hispanic	1 (3.3)	1 (3.3)
American Indian or Alaska Native	1 (3.3)	2 (6.7)
<b>Comorbidities, No. (%)</b>		
Liver Disease	10 (33.3)	7 (23.3)
Seizure Disorder	4 (13.3)	1 (3.3)
Psychiatric Disorder	12 (40)	10 (33.3)
Polysubstance Use	4 (13.3)	1 (3.3)
History of Delirium/Tremens	0	1 (3.3)
<b>Hospital Day 1 Laboratory Values, No. (Mean)</b>		
Baseline ALT	103.4 (range 9-417)	77 (range 8-241)
Baseline AST	135.4 (range 21-611)	107.1 (range 21-520)
CIWA-Ar score	13.6 (range 1-35)	12.7 (range 3-33)
<b>Hospital Course, Mg (Mean)</b>		
Average phenobarbital loading dose	520.5 (range 0-790)	-
Average cumulative phenobarbital dose	862.2 (range 130-4820)	-
Total lorazepam equivalents	18 (range 0-89.7)	15 (range 2-61)

Abbreviations: ED = Emergency Department; ICU = Intensive Care Unit; CIWA = Clinical Institute Withdrawal Assessment for Alcohol; PAWSS = Prediction of Alcohol Withdrawal Severity Scale; No = Number; ALT = Alanine Transaminase; AST = Aspartate Aminotransferase; BZD = Benzodiazepines

Table 2. Clinical Outcomes

Outcome, No. (%)	Phenobarbital Group (n=13)	BZD Group (n=30)	Phenobarbital and BZD Group (n= 17)
Discharge from ED	13 (100)	8 (26.7)	4 (23.5)
ED to ICU admission	0	2 (6.6)	4 (23.5)
ED to floor admission	0	15 (50)	9 (52.9)
Use of restraints	2 (15.4)	5 (16.7)	10 (58.8)
Use of adjunctive agents	1 (7.7)	8 (26.7)	3 (17.6)
Mechanical ventilation	0	0	0

Table 3. Adjunctive Agents Utilized

Medication, No. (%)	Phenobarbital Group (n=13)	BZD Group (n=30)	Phenobarbital and BZD Group (n= 17)
Dexmedetomidine	0	1 (3.3)	2 (11.8)
Propofol	0	0	0
Ketamine	0	0	0
Quetiapine	0	1 (3.3)	3 (17.6)
Haloperidol	0	2 (6.7)	0
Olanzapine	0	3 (10)	0
Droperidol	1 (7.7)	2 (6.7)	0

## Protocol and Education Materials





Figure 1. Prediction of Alcohol Withdrawal Severity Scale (PAWSS)

Target RASS	RASS Description
+4	Combative, violent, danger to staff
+3	Pulls or removes tube(s) or catheters; aggressive
+2	Frequent nonpurposeful movement, fights ventilator
+1	Anxious, apprehensive, but not aggressive
0	Alert and calm
-1	awakens to voice (eye opening/contact) $\geq 10$ sec
-2	light sedation, briefly awakens to voice (eye opening/contact) $\geq 10$ sec
-3	moderate sedation, movement or eye opening. No eye contact
-4	deep sedation, no response to voice, but movement or eye opening to physical stimulation
-5	Unresponsive, no response to voice or physical stimulation

Figure 2. Richmond Agitation Sedation Scale (RASS)

QR Code 1. Oregon Region Phenobarbital for Alcohol Withdrawal in ED Protocol



Protocol

QR Code 2. Nursing Education Video for the Phenobarbital Protocol



Education Video Link

## Discussion

### Patient Population

- At baseline, patients were similar in age, sex, race.
- The mean age was 47 years (range 25-75 years) in the phenobarbital group and 45 years (range 29-70 years) in the BZD group.
- Majority of patients were White or Caucasian.

### Clinical Outcomes

- Patients who were treated with phenobarbital monotherapy did not require further ICU admission.
- Overall rates of further hospital admission were lower in the phenobarbital +/- BZD group compared to the BZD group (43.3% vs. 56.7%, respectively).
- Use of adjunctive agents were lower in the phenobarbital +/- BZD group compared to the BZD group (13.3% vs. 26.7%, respectively).
- Higher use of restraints noted in phenobarbital +/- BZD group compared to BZD (40% vs. 16.7%, respectively).
- Protocol compliance was noted in 60% of patients.

### Study Limitations

- This is a retrospective, non-randomized study.
- Given recent protocol implementation, limitation of time post implementation
- Insufficient sample size required for adequate power to assess the primary and secondary outcomes
- Treatment bias due to patients in phenobarbital group also receiving BZDs and other adjunctive agents.
- Protocol compliance is difficult to assess.
- The baseline severity of alcohol withdrawal varied.

## Preliminary Conclusions

- This study was performed to evaluate the safety and efficacy of phenobarbital for alcohol withdrawal compared to benzodiazepines in patients admitted to the ED at two large tertiary medical centers.
- Doses of phenobarbital and use of adjunctive sedative agents varied in patients due to individual provider prescribing practices.
- Statistical analysis is still underway but descriptive data suggest that phenobarbital may be a safe and effective treatment alternative to BZDs.
- Based on preliminary results, it appears that treatment with phenobarbital may result in reduced hospital admissions compared to standard BZD based regimens.
- Phenobarbital use may result in a reduction in the use of restraints or adjunctive agents.
- Many patients in the phenobarbital group were started on BZDs after receiving a dose of phenobarbital, which could lead to bias of results.
- Data should continue to be collected in a retrospective manner in order to adequately power this study and decrease the level of bias.
- Cost benefit analysis was not conducted in this study but the implications of phenobarbital in preventing further hospital admission, incidence of mechanical ventilation, and use of other adjunctive agents may result in cost savings that may be analyzed in the future.

## References

- Long D, Long B, Koyfman A. The emergency medicine management of severe alcohol withdrawal. Am J Emerg Med. 2017 Jul;35(7):1005-1011. doi: 10.1016/j.ajem.2017.02.002 Epub 2017 Feb 4.
- Wong J, Saver S, Scanlon J, et al. 2020. The ASAM Clinical Practice Guideline on Alcohol Withdrawal Management. J Addict Med. May/Jun 2020;14(35 Suppl 1):1-72. doi: 10.1097/ADM.0000000000000668
- Rosenson J, Clements C, Simon B, et al. Phenobarbital for acute alcohol withdrawal: A prospective randomized double-blind placebo-controlled study. J Emerg Med. 2013;44(3):592-598. doi: 10.1016/j.ajem.2012.07.056
- Tidewell WP, Thomas TL, Posilist JD, Canonica AE, Webber AJ. Treatment of Alcohol Withdrawal Syndrome: Phenobarbital vs CIWA-Ar Protocol. Am J Crit Care. 2018 Nov;27(6):454-460. doi: 10.4037/ajcc.2018.745.
- Parti R, Agne P. Update on Phenobarbital for Alcohol Withdrawal Syndrome in Intensive Care. J Clin Intensive Care Med. 2019; 4: 036-039. doi:10.29328/journal.icim.1001023
- Hammond D, Rowe J, Wong A, et al. 2017. Patient Outcomes Associated With Phenobarbital Use With or Without Benzodiazepines for Alcohol Withdrawal Syndrome: A Systematic Review. Hosp Pharm. 2017 Oct;52(9):607-616. doi: 10.1177/001878171720310. Epub 2017 Jul 17.
- Oks M, Cleven KL, Healy L, et al. The Safety and Utility of Phenobarbital Use for the Treatment of Severe Alcohol Withdrawal Syndrome in the Medical Intensive Care Unit. J Intensive Care Med. 2018;1-7. doi:10.1177/0885066618783947
- Nisavic M, Nejad SH, Isenberg BM, et al. Use of Phenobarbital in Alcohol Withdrawal Management – A Retrospective Comparison Study of Phenobarbital and Benzodiazepines for Acute Alcohol Withdrawal Management in General Medical Patients. Psychosomatics. 2019;60(5):458-467. doi:10.1016/j.psych.2019.02.002
- Hendey GW, Dery RA, Barnes RL, Snowden B, Mentler P. A prospective, randomized, trial of phenobarbital versus benzodiazepines for acute alcohol withdrawal. Am J Emerg Med. 2011;29(4):382-385. doi:10.1016/j.ajem.2009.10.010
- Shah P, Stegner K, Rachid M, Hamir T, Dodd RW. Front-loaded Phenobarbital Dosing Is Associated with a Lower Incidence of Mechanical Ventilation in Patients with Severe Alcohol Withdrawal. 2019;Level 3:A4144-A4144. doi:10.1164/ajrcm-conference